


Parametric Equations: $x = f(t)$, $y = g(t)$

Example: $x = 5t$; $y = t^2 + 6$

Table of Values for Parametric Equations

t	$X(t) = 5t$	$Y(t) = t^2 + 6$
-4	-20	22
-3	-15	15
-2	-10	10
-1	-5	7
0	0	6
1	5	7
2	10	10
3	15	15
4	20	22

$$x = 5t; \quad y = t^2 + 6$$



Parametric Form of the Derivative:

$$\text{a) } \frac{dy}{dx} = \frac{dy/dt}{dx/dt}$$

$$\text{b) } \frac{d^2 y}{dx^2} = \frac{\frac{d}{dt} \left[\frac{dy}{dx} \right]}{dx/dt}$$

Example 1:

Parametric Equations: $x = 4t^2$, $y = 4 - 3t$

Find $y' = \frac{dy}{dx}$ = Derivative of y with respect to x .

We can find $\frac{dy}{dx}$ by using the Direct Method or Chain Rule Method.

Direct Method:

$$x = 4t^2 \Rightarrow t^2 = \frac{x}{4} \Rightarrow \sqrt{t^2} = \pm \sqrt{\frac{x}{4}} \Rightarrow t = \pm \frac{\sqrt{x}}{2}$$

$$y = 4 - 3t \Rightarrow y = 4 - 3\left(\pm \frac{\sqrt{x}}{2}\right) \Rightarrow y = 4 \pm \frac{3\sqrt{x}}{2} \Rightarrow y = 4 \pm \frac{3}{2}x^{1/2}$$

$$y' = \pm \frac{3}{2} \left(\frac{1}{2} x^{-1/2} \right) = \pm \frac{3}{4} x^{-1/2} = \pm \frac{3}{4x^{1/2}} = \pm \frac{3}{4\sqrt{x}}$$

Parametric Equations: $x = 4t^2$, $y = 4 - 3t$

Chain Rule Method:

$$\text{Note: } x = 4t^2 \Rightarrow t^2 = \frac{x}{4} \Rightarrow \sqrt{t^2} = \pm \sqrt{\frac{x}{4}} \Rightarrow t = \pm \frac{\sqrt{x}}{2}$$

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-3}{8t} = \frac{-3}{8\left(\pm \frac{\sqrt{x}}{2}\right)} = \frac{-3}{\pm 4\sqrt{x}} = \pm \frac{3}{4\sqrt{x}}$$

Parametric Equations: $x = 4t^2$, $y = 4 - 3t$

We can also find y'' by using the Chain Rule Method:

$$\frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-3}{8t} = \frac{-3}{8\left(\pm\frac{\sqrt{x}}{2}\right)} = \frac{-3}{\pm 4\sqrt{x}} = \pm \frac{3}{4\sqrt{x}}$$

$$\begin{aligned} y'' &= \frac{d^2 y}{dx^2} = \frac{\frac{d}{dt} \left[\frac{dy}{dx} \right]}{dx/dt} = \frac{\frac{d}{dt} \left[\frac{-3}{8t} \right]}{8t} = \frac{\frac{d}{dt} \left[-\frac{3}{8} t^{-1} \right]}{8t} = \frac{\frac{3}{8} t^{-2}}{8t} = \frac{3}{64t^3} \\ &= \frac{3}{64 \left(\pm \frac{3}{4\sqrt{x}} \right)^3} \end{aligned}$$

Example 2:

Parametric Equations: $x = 4\sqrt{t}$, $y = 2t$

Find $y' = \frac{dy}{dx}$

Note : $x = 4\sqrt{t} = 4t^{1/2}$

Find $y' = \frac{dy}{dx}$ by using the Direct Method.

$$x = 4\sqrt{t} \Rightarrow (x)^2 = (4\sqrt{t})^2 \Rightarrow x^2 = 16t \Rightarrow t = \frac{x^2}{16}$$

$$y = 2t \Rightarrow y = 2\left(\frac{x^2}{16}\right) \Rightarrow y = \frac{1}{8}x^2$$

$$y' = \frac{1}{8}(2x) = \frac{1}{4}x$$

Example 2:

Parametric Equations: $x = 4\sqrt{t}$, $y = 2t$

Find $y' = \frac{dy}{dx}$ and y'' by using the Chain Rule Method.

$$x = 4\sqrt{t} \Rightarrow (x)^2 = (4\sqrt{t})^2 \Rightarrow x^2 = 16t \Rightarrow t = \frac{x^2}{16}$$

$$y' = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2}{2t^{-1/2}} = \frac{1}{t^{-1/2}} = t^{1/2} = \left(\frac{x^2}{16}\right)^{1/2} = \frac{x}{4}$$

$$y'' = \frac{d^2 y}{dx^2} = \frac{\frac{d}{dt} \left[\frac{dy}{dx} \right]}{dx/dt} = \frac{\frac{d}{dt} \left[t^{1/2} \right]}{2t^{-1/2}} = \frac{\frac{1}{2} t^{-1/2}}{2t^{-1/2}} = \frac{1}{2} = \frac{1}{4}$$

Example 3:

Parametric Equations: $x = 2t$, $y = 5 - t^2$

Find the equation of the tangent line passing through (2,4).

At the point (2,4), $x = 2$ and $y = 4$. To find t at this point, we can use $x = 2t$.

$$x = 2t \quad \Rightarrow \quad 2 = 2t \quad \Rightarrow \quad t = 1$$

To find the slope of the tangent line, we need to find y' .

$$y' = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{-2t}{2} = -t$$

Example 3: (con't)


$$\text{Slope of tangent line} = y' = \frac{dy}{dx} = -t = -(1) = -1$$

Equation of tangent line at $(2,4)$:

$$y - y_1 = m(x - x_1)$$

$$y - 4 = -1(x - 2)$$

Parametric Equations: $x = 2t$, $y = 5 - t^2$



Example 4:

Parametric Equations: $x = 2 \cos t$, $y = 2 \sin t$ $0 \leq t \leq 2\pi$

Find the equation of the tangent line at the point $(0,2)$.

At the point $(0,2)$, we can find t as follows:

$$x = 2 \cos t \quad \Leftrightarrow \quad 0 = 2 \cos t \quad \Leftrightarrow \quad \cos t = 0 \quad \Leftrightarrow \quad t = \pi / 2$$

To find the slope of the tangent line, we need to find y' .

$$y' = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{2 \cos t}{-2 \sin t} = -\frac{\cos t}{\sin t}$$

Example 4 (con't):

$$\text{Slope of tangent line} = \frac{dy}{dx} = -\frac{\cos t}{\sin t} = -\frac{\cos(\pi/2)}{\sin(\pi/2)} = \frac{0}{1} = 0$$

Equation of tangent line at $(0,2)$:

$$y - y_1 = m(x - x_1)$$

$$y - 2 = 0(x - 0)$$

$$y - 2 = 0$$

Calculator Input:

Parametric Equations [Video](#) on/off

Tmin = 0

Tmax = 2π

Tstepsize = 0.01

Example

Submit

clear

Table of Values

Tracing Graph

$$X(t) = 2\cos(t)$$


1

on/off

$$Y(t) = 2\sin(t)$$

1

$$x = 2 \cos t, \quad y = 2 \sin t$$



Example 5:

Parametric Equations: $x = 4t + t^2$, $y = 2t^{3/2}$

Find equation of the tangent line at $(5,2)$.

$$y' = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{3t^{1/2}}{4 + 2t}$$

At the point $(5,2)$:

$$y = 2t^{3/2} \quad \Leftrightarrow \quad 2 = 2t^{3/2} \quad \Leftrightarrow \quad t^{3/2} = 1 \quad \Leftrightarrow \quad t = 1$$

Example 5 (con't):

Parametric Equations: $x = 4t + t^2$, $y = 2t^{3/2}$

$$\text{Slope of tangent line} = \frac{dy}{dx} = \frac{3t^{1/2}}{4 + 2t} = \frac{3(1)^{1/2}}{4 + 2(1)} = \frac{3}{6} = \frac{1}{2}$$

Equation of tangent line at $(5, 2)$:

$$y - y_1 = m(x - x_1)$$

$$y - 2 = \frac{1}{2}(x - 5)$$


Calculator Input:

Parametric Equations [Video](#) on/off

Tmin = Tmax = Tstepsize =

X(t) = on/off

Y(t) =



Example 6:


Parametric Equations: $x = 6t + 2$, $y = 3t$ $0 \leq t \leq 2$

$$s = \text{Arc Length} = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt :$$

$$\frac{dx}{dt} = 6 \qquad \frac{dy}{dt} = 3$$

$$\begin{aligned} s &= \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_0^2 \sqrt{(6)^2 + (3)^2} dt = \sqrt{45}t \Big|_0^2 \\ &= 2\sqrt{45} - 0 = 2\sqrt{45} \end{aligned}$$

$$x = 6t + 2, \quad y = 3t \quad 0 \leq t \leq 2$$



Example 7:


Parametric Equations: $x = t^2 + 3$, $y = t^3 - 2$ $0 \leq t \leq 2$

$$s = \text{Arc Length} = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt :$$

$$\frac{dx}{dt} = 2t \qquad \frac{dy}{dt} = 3t^2$$

$$s = \int_0^2 \sqrt{(2t)^2 + (3t^2)^2} dt = \int_0^2 \sqrt{4t^2 + 9t^4} dt = 9.0734$$

$$x = t^2 + 3, \quad y = t^3 - 2 \quad 0 \leq t \leq 2$$



Example 8:

Parametric Equations: $x = 4 \cos t + 1$, $y = 3 \sin t$ $0 \leq t \leq \pi / 4$

$$s = \text{Arc Length} = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt :$$

$$\frac{dx}{dt} = 4(-\sin t) = -4 \sin t; \quad \frac{dy}{dt} = 3 \cos t$$

$$s = \int_0^{\pi/4} \sqrt{(-4 \sin t)^2 + (3 \cos t)^2} dt = \int_0^{\pi/4} \sqrt{16(\sin t)^2 + 9(\cos t)^2} dt$$
$$= 2.51378$$

Calculator Input:


Parametric Equations [Video](#) on/off

Tmin = Tmax = Tstepsize =

X(t) = on/off

Y(t) =

$$x = 4 \cos t + 1, \quad y = 3 \sin t \quad 0 \leq t \leq \pi/4$$



Find $\int_0^{\pi/4} \sqrt{16(\sin t)^2 + 9(\cos t)^2} dt$ by using Trapezoidal Rule.

$$\int_0^{\pi/4} \sqrt{16(\sin t)^2 + 9(\cos t)^2} dt = 2.51378$$

Input for finding definite integral

Trapezoidal Rule

Input f(x) = $\sqrt{16(\sin(x))^2 + 9(\cos(x))^2}$ Find Area

Input value of a = **0**

Input value of b = **$\pi/4$**

Input value of n = number of subintervals = **5000**

Example 9:

Parametric Equations: $x = 2e^t$, $y = e^{3t}$ $0 \leq t \leq 1$

$$s = \text{Arc Length} = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt :$$

$$\frac{dx}{dt} = 2e^t \quad \frac{dy}{dt} = e^{3t} (3) = 3e^{3t}$$

$$s = \int_0^1 \sqrt{(2e^t)^2 + (3e^{3t})^2} dt = \int_0^1 \sqrt{4e^{2t} + 9e^{6t}} dt = 19.4937$$

Calculator Input

Parametric Equations [Video](#) on/off

Tmin = 0

Tmax = 1

Tstepsize = 0.01

Example

Submit

clear

Table of Values

Tracing Graph

$$X(t) = 2e^{(t)}$$


1

on/off

$$Y(t) = e^{(3t)}$$

1

$$x = 2e^t, \quad y = e^{3t} \quad 0 \leq t \leq 1$$



Find $\int_0^1 \sqrt{4e^{2t} + 9e^{6t}} dt$ by using Trapezoidal Rule.

$$\int_0^1 \sqrt{4e^{2t} + 9e^{6t}} dt = 19.4937$$

Input for finding definite integral

Trapezoidal Rule

Input $f(x) = \sqrt{(4e^{(2x)} + 9e^{(6x)})}$ Find Area

Input value of a = **0**

Input value of b = **1**

Input value of n = number of subintervals = **5000**