

Vectors in the Plane


Scalar Quantities: area, volume, height, temperature, time,...

Vector Quantity is a quantity of something which possesses both magnitude and direction.

Examples Vector Quantities: force, velocity, acceleration, ...


Representing Vector with directed line segment:

A car's velocity of 50 miles per hour, going northeast.




$$\text{Length or Magnitude of Vector} = \|\overrightarrow{PQ}\| = 30$$

A car's acceleration of 20 miles per hour, going east.



$$\text{Length or Magnitude of Vector} = \|\overrightarrow{PQ}\| = 20$$

Find the length of vector \overrightarrow{SR} .




Length or Magnitude of Vector = $\|\overrightarrow{SR}\|$

= distance between R and S

$$= \sqrt{(10 - 2)^2 + (10 - 2)^2} = 11.3137$$

Find the length of vector \overrightarrow{SR} .




$$\text{Length or Magnitude of Vector} = \|\overrightarrow{SR}\|$$

$$= \text{distance between R and S}$$

$$= \sqrt{(10 - 4)^2 + (-8 - -2)^2} = 8.4852813$$


Vector in Standard Position: Vector with initial point at origin.



Components of vector \mathbf{v} are v_1 and v_2 .

$$\text{Magnitude of } \mathbf{v} = \|\mathbf{v}\| = \|\langle v_1, v_2 \rangle\| = \sqrt{(10)^2 + (10)^2} = \sqrt{200}$$

Vector in Non-Standard Position:



Components of vector \mathbf{v} are v_1 and v_2 :

$$\langle v_1, v_2 \rangle = \langle q_1 - p_1, q_2 - p_2 \rangle = \langle 10 - 2, 7 - (-1) \rangle = \langle 8, 8 \rangle$$

Magnitude of \mathbf{v} :

$$\text{Magnitude of } \mathbf{v} = \|\mathbf{v}\| = \|\langle v_1, v_2 \rangle\| = \sqrt{(8)^2 + (8)^2} = \sqrt{128}$$

Vector Operations:

$$\text{Let } \mathbf{u} = \langle u_1, u_2 \rangle \text{ and } \mathbf{v} = \langle v_1, v_2 \rangle$$

$$\text{a) } \mathbf{u} + \mathbf{v} = \langle u_1, u_2 \rangle + \langle v_1, v_2 \rangle = \langle u_1 + v_1, u_2 + v_2 \rangle$$

$$\text{b) } -\mathbf{u} = \langle -u_1, -u_2 \rangle; \quad -\mathbf{v} = \langle -v_1, -v_2 \rangle$$

$$\text{c) } \mathbf{u} - \mathbf{v} = \mathbf{u} + (-\mathbf{v}) = \langle u_1, u_2 \rangle + \langle -v_1, -v_2 \rangle = \langle u_1 + -v_1, u_2 + -v_2 \rangle$$

d) Scalar Multiple:

$$b\mathbf{u} = b\langle u_1, u_2 \rangle = \langle bu_1, bu_2 \rangle$$

$$c\mathbf{v} = c\langle v_1, v_2 \rangle = \langle cv_1, cv_2 \rangle$$

Examples:

$$\text{Let } \mathbf{u} = \langle 3, 7 \rangle \text{ and } \mathbf{v} = \langle 1, 8 \rangle$$

$$\text{a) } \mathbf{u} + \mathbf{v} = \langle 3, 7 \rangle + \langle 1, 8 \rangle = \langle 4, 15 \rangle$$

$$\text{b) } -\mathbf{u} = \langle -3, -7 \rangle; \quad -\mathbf{v} = \langle -1, -8 \rangle$$

$$\text{c) } \mathbf{u} - \mathbf{v} = \mathbf{u} + (-\mathbf{v}) = \langle 3, 7 \rangle + \langle -1, -8 \rangle = \langle 2, -1 \rangle$$

d) Scalar Multiple:

$$9\mathbf{u} = 9\langle 3, 7 \rangle = \langle 27, 63 \rangle$$

$$-3\mathbf{v} = -3\langle 1, 8 \rangle = \langle -3, -24 \rangle$$

$$e) 3\mathbf{u} + 5\mathbf{v} = 3\langle 3, 7 \rangle + 5\langle 1, 8 \rangle = \langle 9, 21 \rangle + \langle 5, 40 \rangle = \langle 14, 61 \rangle$$

$$f) 8\mathbf{u} - 6\mathbf{v} = 8\mathbf{u} + (-6\mathbf{v}) = 8\langle 3, 7 \rangle + (-6)\langle 1, 8 \rangle \\ = \langle 24, 56 \rangle + \langle -6, -48 \rangle = \langle 18, 8 \rangle$$

Unit Vector = vector with length of 1.


Let \mathbf{u} be a unit vector in the direction of \mathbf{v} :

$$\text{Then } \mathbf{u} = \frac{\mathbf{v}}{\|\mathbf{v}\|} = \frac{1}{\|\mathbf{v}\|} \mathbf{v} \quad \text{Note: } \mathbf{v} \text{ is a vector; } \|\mathbf{v}\| \text{ is a scalar.}$$

Show that $\|\mathbf{u}\| = 1$.

$$\|\mathbf{u}\| = \left\| \frac{1}{\|\mathbf{v}\|} \mathbf{v} \right\| = \frac{1}{\|\mathbf{v}\|} \|\mathbf{v}\| = \frac{\|\mathbf{v}\|}{\|\mathbf{v}\|} = 1$$

Example:



Components of vector \mathbf{v} are v_1 and v_2 :


$$\langle v_1, v_2 \rangle = \langle q_1 - p_1, q_2 - p_2 \rangle = \langle 10 - 2, 7 - (-1) \rangle = \langle 8, 8 \rangle$$

Magnitude of \mathbf{v} :

$$\text{Magnitude of } \mathbf{v} = \|\mathbf{v}\| = \|\langle v_1, v_2 \rangle\| = \sqrt{(8)^2 + (8)^2} = \sqrt{128}$$

$$\mathbf{u} = \frac{\mathbf{1}}{\|\mathbf{v}\|} \mathbf{v} = \frac{1}{\sqrt{128}} \langle 8, 8 \rangle = \left\langle \frac{8}{\sqrt{128}}, \frac{8}{\sqrt{128}} \right\rangle$$

$$\|\mathbf{u}\| = \left\| \left\langle \frac{8}{\sqrt{128}}, \frac{8}{\sqrt{128}} \right\rangle \right\| = \sqrt{\left(\frac{8}{\sqrt{128}}\right)^2 + \left(\frac{8}{\sqrt{128}}\right)^2} = 1$$



\mathbf{u} has length of 1 and \mathbf{u} is the same direction as \mathbf{v} .

Standard Unit Vectors:

$$\mathbf{i} = \langle 1, 0 \rangle \quad \text{and} \quad \mathbf{j} = \langle 0, 1 \rangle$$

$$\text{Let } \mathbf{v} = \langle v_1, v_2 \rangle$$

Show that $\mathbf{v} = v_1\mathbf{i} + v_2\mathbf{j}$:

$$v_1\mathbf{i} + v_2\mathbf{j} = v_1\langle 1, 0 \rangle + v_2\langle 0, 1 \rangle = \langle v_1, 0 \rangle + \langle 0, v_2 \rangle = \langle v_1, v_2 \rangle = \mathbf{v}$$

Example: Let $\mathbf{v} = \langle 2, 7 \rangle$

Then \mathbf{v} can be expressed with unit vectors as follows;

$$\mathbf{v} = 2\mathbf{i} + 7\mathbf{j}$$

Example: Let $\mathbf{u} = \langle -2, 5 \rangle$

Then \mathbf{u} can be expressed with unit vectors as follows;

$$\mathbf{u} = -2\mathbf{i} + 5\mathbf{j}$$

Example: Let $\mathbf{u} = 7\mathbf{i} + 9\mathbf{j}$

Find magnitude of \mathbf{u} .

Note: $\mathbf{u} = 7\mathbf{i} + 9\mathbf{j} = \langle 7, 9 \rangle$

$$\|\mathbf{u}\| = \sqrt{(7)^2 + (9)^2} = \sqrt{112}$$

Example: Let $\mathbf{u} = \langle 7, 9 \rangle$ and $\mathbf{v} = \langle 1, 2 \rangle$

$$\|\mathbf{u}\| = \sqrt{(7)^2 + (9)^2} = \sqrt{130}$$

$$\|\mathbf{v}\| = \sqrt{(1)^2 + (2)^2} = \sqrt{5}$$

$$\mathbf{u} + \mathbf{v} = \langle 8, 11 \rangle$$

$$\|\mathbf{u} + \mathbf{v}\| = \sqrt{(8)^2 + (11)^2} = \sqrt{185}$$

Note: $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$ (or $\sqrt{185} \leq \sqrt{130} + \sqrt{5}$)

$$\left\| \frac{\mathbf{u}}{\|\mathbf{u}\|} \right\| = \left\| \frac{1}{\|\mathbf{u}\|} \cdot \mathbf{u} \right\| = \frac{1}{\|\mathbf{u}\|} \|\mathbf{u}\| = \frac{1}{\sqrt{130}} \sqrt{130} = 1$$

$$\left\| \frac{\mathbf{v}}{\|\mathbf{v}\|} \right\| = \left\| \frac{1}{\|\mathbf{v}\|} \cdot \mathbf{v} \right\| = \frac{1}{\|\mathbf{v}\|} \|\mathbf{v}\| = \frac{1}{\sqrt{5}} \sqrt{5} = 1$$

$$\left\| \frac{\mathbf{u} + \mathbf{v}}{\|\mathbf{u} + \mathbf{v}\|} \right\| = \left\| \frac{1}{\|\mathbf{u} + \mathbf{v}\|} \cdot (\mathbf{u} + \mathbf{v}) \right\| = \frac{1}{\|\mathbf{u} + \mathbf{v}\|} \|\mathbf{u} + \mathbf{v}\| = \frac{1}{\sqrt{185}} \sqrt{185} = 1$$

Find vector \mathbf{v} with given magnitude that has same direction as \mathbf{u} :

Let $\|\mathbf{v}\| = 3$ and $\mathbf{u} = \langle 4, 7 \rangle$

Note: $\|\mathbf{u}\| = \sqrt{4^2 + 7^2} = \sqrt{65}$

Let \mathbf{w} be the unit vector in the same direction as \mathbf{u}

$$\mathbf{w} = \frac{1}{\|\mathbf{u}\|} \mathbf{u} = \frac{1}{\sqrt{65}} \langle 4, 7 \rangle = \left\langle \frac{4}{\sqrt{65}}, \frac{7}{\sqrt{65}} \right\rangle$$

Hence, \mathbf{w} and \mathbf{u} are in the same direction.

Then, $3\mathbf{w}$ would be a vector of magnitude 3 and has same direction as \mathbf{w} and \mathbf{u} .

$$3\mathbf{w} = 3 \left\langle \frac{4}{\sqrt{65}}, \frac{7}{\sqrt{65}} \right\rangle = \left\langle \frac{12}{\sqrt{65}}, \frac{21}{\sqrt{65}} \right\rangle$$

$$\|3\mathbf{w}\| = \left\| \left\langle \frac{12}{\sqrt{65}}, \frac{21}{\sqrt{65}} \right\rangle \right\| = \sqrt{\left(\frac{12}{\sqrt{65}} \right)^2 + \left(\frac{21}{\sqrt{65}} \right)^2} = 3$$

Example of Linear Combination

Let $\mathbf{u} = \langle 1, 2 \rangle$, $\mathbf{v} = \langle 7, 8 \rangle$, and $\mathbf{w} = \langle -3, 4 \rangle$.

Find scalars a and b such that $\mathbf{v} = a\mathbf{u} + b\mathbf{w}$.

Let $\mathbf{u} = \langle 1, 2 \rangle$, $\mathbf{v} = \langle 7, 8 \rangle$, and $\mathbf{w} = \langle -3, 4 \rangle$.

Note: $\mathbf{u} = \langle 1, 2 \rangle = \mathbf{i} + 2\mathbf{j}$ and $\mathbf{w} = \langle -3, 4 \rangle = -3\mathbf{i} + 4\mathbf{j}$.

$$\mathbf{v} = a\mathbf{u} + b\mathbf{w} = a(\mathbf{i} + 2\mathbf{j}) + b(-3\mathbf{i} + 4\mathbf{j})$$

$$= a\mathbf{i} + 2a\mathbf{j} - 3b\mathbf{i} + 4b\mathbf{j}$$

$$= a\mathbf{i} - 3b\mathbf{i} + 2a\mathbf{j} + 4b\mathbf{j}$$

$$= (a - 3b)\mathbf{i} + (2a + 4b)\mathbf{j}$$

$$\langle 7, 8 \rangle = \langle a - 3b, 2a + 4b \rangle$$

$$a - 3b = 7 \quad (1)$$

$$2a + 4b = 8 \quad (2)$$

Solving system of two equations:

$$(1) \quad a = 3b + 7$$

$$(2) \quad 2(3b + 7) + 4b = 8$$

$$10b + 14 = 8$$

$$b = -6/10 = -3/5$$

$$a = 3(-3/5) + 7 = 26/5$$

Summary:

$$\mathbf{v} = a\mathbf{u} + b\mathbf{w}$$

$$\langle 7, 8 \rangle = \frac{-3}{5} \langle 1, 2 \rangle + \frac{26}{5} \langle -3, 4 \rangle$$