


$$f(x) = 2 \ln x$$

a) Domain of  $f(x) = \{x : x > 0\}$


b) Range of  $f(x) = (-\infty, \infty)$



$$f(x) = \ln(x + 4)$$

a) Domain of  $f(x) = \{x : x + 4 > 0\} = \{x : x > -4\}$

b) Range of  $f(x) = (-\infty, \infty)$



## Properties of Logarithm:

Product Rule:  $\ln(A \cdot B) = \ln A + \ln B$

Quotient Rule:  $\ln(A / B) = \ln A - \ln B$

Power Rule:  $\ln(x^p) = p \ln x$

Product Rule:  $\ln(A \cdot B) = \ln A + \ln B$

Expand  $\ln(4 \cdot x)$ .

$$\ln(4 \cdot x) = \ln 4 + \ln x$$

Quotient Rule:  $\ln(A/B) = \ln A - \ln B$

Expand  $\ln \frac{x}{5}$ .

$$\ln \frac{x}{5} = \ln x - \ln 5$$

Expand  $\ln \frac{ab}{z}$ .

Properties of Logarithm:

Product Rule:  $\ln(A \cdot B) = \ln A + \ln B$

Quotient Rule:  $\ln(A / B) = \ln A - \ln B$

Power Rule:  $\ln(x^p) = p \ln x$

$$\ln \frac{ab}{z} = \ln(ab) - \ln(z) = \ln(a) + \ln(b) - \ln(z)$$



Expand  $\ln\left(x^2\sqrt{x^4+5}\right)$ .

Properties of Logarithm:

Product Rule:  $\ln(A \cdot B) = \ln A + \ln B$

Quotient Rule:  $\ln(A / B) = \ln A - \ln B$

Power Rule:  $\ln(x^p) = p \ln x$

$$\begin{aligned}\ln\left(x^2\sqrt{x^4+5}\right) &= \ln\left(x^2\right) + \ln\left(\sqrt{x^4+5}\right) \\ &= \ln\left(x^2\right) + \ln\left(\left[x^4+5\right]^{1/2}\right) = 2\ln(x) + \frac{1}{2}\ln\left(x^4+5\right)\end{aligned}$$

$$\text{Expand } \ln\left(\sqrt{\frac{2x-4}{x}}\right) = \ln\left(\frac{2x-4}{x}\right)^{1/2}$$

Properties of Logarithm:

$$\text{Product Rule: } \ln(A \cdot B) = \ln A + \ln B$$

$$\text{Quotient Rule: } \ln(A / B) = \ln A - \ln B$$

$$\text{Power Rule: } \ln(x^p) = p \ln x$$

$$\begin{aligned} \ln\left(\sqrt{\frac{2x-4}{x}}\right) &= \ln\left(\frac{2x-4}{x}\right)^{1/2} = \frac{1}{2} \ln\left(\frac{2x-4}{x}\right) \\ &= \frac{1}{2} [\ln(2x-4) - \ln x] = \frac{1}{2} \ln(2x-4) - \frac{1}{2} \ln(x) \end{aligned}$$

Combine:  $\ln(x - 4) - \ln(x + 5)$ .

Properties of Logarithm:

Product Rule:  $\ln(A \cdot B) = \ln A + \ln B$

Quotient Rule:  $\ln(A / B) = \ln A - \ln B$

Power Rule:  $\ln(x^p) = p \ln x$

$$\ln(x - 4) - \ln(x + 5) = \ln\left(\frac{x - 4}{x + 5}\right)$$

Combine:  $2\ln(x^2 + 5) + \ln x - \ln(x - 11)$

Properties of Logarithm:

Product Rule:  $\ln(A \cdot B) = \ln A + \ln B$

Quotient Rule:  $\ln(A / B) = \ln A - \ln B$

Power Rule:  $\ln(x^p) = p \ln x$

$$2\ln(x^2 + 5) + \ln x - \ln(x - 11) = \ln(x^2 + 5)^2 + \ln x - \ln(x - 11)$$

$$= \ln\left(\frac{(x^2 + 5)^2 \cdot x}{x - 11}\right)$$

$$\text{Combine: } 2\ln 7 - \frac{1}{3}\ln(x^4 + 1).$$

Properties of Logarithm:

$$\text{Product Rule: } \ln(A \cdot B) = \ln A + \ln B$$

$$\text{Quotient Rule: } \ln(A / B) = \ln A - \ln B$$

$$\text{Power Rule: } \ln(x^p) = p \ln x$$

$$2\ln 7 - \frac{1}{3}\ln(x^4 + 1) = \ln 7^2 - \ln(x^4 + 1)^{1/3}$$

$$= \ln 49 - \ln(x^4 + 1)^{1/3} = \ln\left(\frac{49}{(x^4 + 1)^{1/3}}\right) = \ln\left(\frac{49}{\sqrt[3]{x^4 + 1}}\right)$$

## Derivative of function with logarithm.

Chain Rule:  $f(x) = \text{Ln}(\text{expression})$

$$f'(x) = \frac{1}{\text{expression}} \cdot D_x(\text{expression})$$

$$f(x) = \ln(7x)$$

$$f'(x) = \frac{1}{7x} \cdot D_x(7x) = \frac{1}{7x} \cdot 7 = \frac{1}{x}$$

Note: $D_x(\ln x) = \frac{1}{x}$
----------------------------------

## Review:

$(\ln x)^3$  is not the same as  $\ln x^3 = 3 \ln x$

$(\ln x)^4$  is not the same as  $\ln x^4 = 4 \ln x$

$(\ln x)^5$  is not the same as  $\ln x^5 = 5 \ln x$

$(\ln x)^6$  is not the same as  $\ln x^6 = 6 \ln x$

$(\ln x)^7$  is not the same as  $\ln x^7 = 7 \ln x$

$(\ln x)^8$  is not the same as  $\ln x^8 = 8 \ln x$

Let  $f(x) = (\ln x)^3$ . Find  $f'(x) = \underline{\hspace{10em} ? \hspace{10em}}$

Power Rule for derivative:  $f(x) = (\text{base})^n$

$$f'(x) = n(\text{base})^{n-1} \cdot D_x(\text{base})$$

$$f(x) = (\ln x)^3$$

Note:  $(\ln x)^3$  is not the same as  $\ln x^3 = 3 \ln x$

$$f'(x) = 3(\ln x)^2 \cdot D_x(\ln x) = 3(\ln x)^2 \cdot \left(\frac{1}{x}\right)$$

$$= \frac{3(\ln x)^2}{x}$$



Let  $f(x) = \ln \left[ 2x \cdot \sqrt{x^2 - 14} \right]$ . Find  $f'(x) = \underline{\hspace{2cm}}$

Chain Rule:  $f(x) = \ln(\text{expression})$

$$\Rightarrow f'(x) = \frac{1}{\text{expression}} \cdot D_x(\text{expression})$$

$$\begin{aligned} f(x) &= \ln \left[ 2x \cdot \sqrt{x^2 - 14} \right] = \ln [2x] + \ln \left[ \sqrt{x^2 - 14} \right] \\ &= \ln [2x] + \ln \left[ (x^2 - 14)^{1/2} \right] = \ln [2x] + \frac{1}{2} \ln [x^2 - 14] \end{aligned}$$

$$f'(x) = \frac{1}{2x} D_x(2x) + \frac{1}{2} \cdot \left[ \frac{1}{x^2 - 14} D_x(x^2 - 14) \right]$$

$$f'(x) = \frac{1}{2x} (2) + \frac{1}{2} \cdot \left[ \frac{1}{x^2 - 14} (2x) \right] = \frac{1}{x} + \frac{x}{x^2 - 14}$$

Let  $f(x) = \ln\left(\frac{\ln x}{x^4}\right)$ . Find  $f'(x) = \underline{\hspace{2cm}}$

Chain Rule:  $f(x) = \ln(\text{expression})$

$$\Rightarrow f'(x) = \frac{1}{\text{expression}} \cdot D_x(\text{expression})$$

$$f(x) = \ln\left(\frac{\ln x}{x^4}\right) = \ln(\ln x) - \ln(x^4) = \ln(\ln x) - 4\ln(x)$$

$$\begin{aligned} f'(x) &= \frac{1}{\ln x} D_x(\ln x) - 4 \left[ \frac{1}{x} \right] = \frac{1}{\ln x} \left( \frac{1}{x} \right) - 4 \left[ \frac{1}{x} \right] \\ &= \frac{1}{x \ln x} - \frac{4}{x} \end{aligned}$$

Let  $y = \ln \sqrt{\frac{x+2}{x-4}}$ . Find  $y' =$  \_\_\_\_\_

$$\text{Note: } \ln \sqrt{\frac{x+2}{x-4}} = \ln \left( \frac{x+2}{x-4} \right)^{1/2} = \frac{1}{2} [\ln(x+2) - \ln(x-4)]$$

$$y' = \frac{1}{2} \left[ \frac{1}{x+2} D_x(x+2) - \frac{1}{x-4} D_x(x-4) \right]$$

$$y' = \frac{1}{2} \left[ \frac{1}{x+2} (1) - \frac{1}{x-4} (1) \right] = \frac{1}{2} \left[ \frac{1}{x+2} - \frac{1}{x-4} \right]$$

Let  $f(x) = \ln x^2 + 2$ . Find Equation of Tangent Line at (1,2)

Note:  $f(x) = \ln x^2 + 2 = 2 \ln x + 2$


a) Find  $f'(x) = 2 \left( \frac{1}{x} \right) = \frac{2}{x}$

b) Slope of Tangent Line  $= f'(1) = \frac{2}{x} = \frac{2}{1} = 2$

c) Equation of Tangent Line at (1,2):  $y - y_1 = m(x - x_1)$

$$y - 2 = 2(x - 1)$$

$$y = 2(x - 1) + 2$$



Let  $f(x) = \ln \sqrt{1 + \sin x}$ . Find Tangent Line at  $(\pi, 0)$

Note:  $\ln \sqrt{1 + \sin x} = \ln (1 + \sin x)^{1/2} = \frac{1}{2} \ln (1 + \sin x)$

a) Find  $f'(x) = \frac{1}{2} \left[ \frac{1}{1 + \sin x} \right] D_x (1 + \sin x) = \frac{1}{2} \left[ \frac{1}{1 + \sin x} \right] (\cos x)$


b) Slope of Tangent Line  $= f'(\pi) = \frac{1}{2} \left[ \frac{1}{1 + \sin \pi} \right] (\cos \pi) = -\frac{1}{2}$

Note:  $\sin \pi = 0$ ;  $\cos \pi = -1$

c) Equation of Tangent Line:  $y - y_1 = m(x - x_1)$

$$y - 0 = -\frac{1}{2}(x - \pi)$$

$$y = -\frac{1}{2}(x - \pi)$$



Let  $f(x) = 4x^2 - \ln x$ . Find Extremum.

a) Find  $f'(x) = 8x - \frac{1}{x}$

b) Finding relative extremum:

$$\text{Set } 8x - \frac{1}{x} = 0$$

$$8x \cdot x - \frac{1}{x} \cdot x = 0 \cdot x \quad \text{Multiply each term by } x$$


$$8x^2 - 1 = 0$$

$$8x^2 = 1 \quad \Rightarrow \quad x^2 = 1/8 \quad \Rightarrow \quad x = \pm\sqrt{1/8}$$

Therefore,  $f(x) = 4x^2 - \ln x$  has minimum at  $x = \sqrt{1/8}$ .

Minimum at  $x = \sqrt{1/8}$  ;  $y = 1.539720770839918$




$$f(x) = 4x^2 - \ln x.$$

Minimum at  $x = \sqrt{1/8}$  ;  $y = 1.539720770839918$